

A3.3a. ESTUDIO DE LA PERDIDA DE RIQUEZA ESPECIFICA EN LA FLORA DE LAS ZONAS DE RECOLECCION PAM EN EXTREMADURA

ACTION 1.- CONSERVACION PAM

AUTORES: Francisco M. Vázquez Pardo, David
García Alonso, Francisco Márquez García y María
José Guerra Barrena.

PROJECT ACRONYM

COOP4PAM

PROJECT TITLE

Cooperar para crescer no setor das plantas aromáticas e medicinais

PROJECT CODE

0665_COOP4PAM_4_P

Deliverable number and name: ESTUDIO DE LA PERDIDA DE RIQUEZA ESPECIFICA EN LA FLORA DE LAS ZONAS DE RECOLECCION PAM EN EXTREMADURA

Work Package: ACTION 1

Author: CICYTEX

DOCUMENT HISTORY

Version	Date of issue	Content and changes	Edited by
	25-X-2020		<i>Francisco M^a Vázquez Pardo David García Alonso Francisco Márquez García María José Guerra Barrena</i>

Colocar texto (legendagem)

The COOP4PAM project is implemented by the following partners:



CONTENT

1	INTRODUCTION	06
2	DATA APPROACH	07
3	BIODIVERSITY CHANGE	08
4	FIGURES	11
5	BIBLIOGRAPHY	13

FIGURES

TABLES

TITLE

Notes of changes in biodiversity in the exploited populations of *Cistus ladanifer* L., (Cistaceae) from SW Iberian Peninsula.

Francisco Márquez García ¹ (Orcid: 0000-0002-4420-2822)

David García Alonso ¹ (Orcid: 0000-0002-9593-6164)

Francisco María Vázquez Pardo ¹ (Orcid: 0000-0001-7826-8814)

Address: Unidad Biodiversidad Vegetal/ CICYTEX-La Orden/ Ctra Madrid-Lisboa km 372/ 06187 Guadajira (Badajoz, Spain)

Contact: frvazquez50@hotmail.com

ABSTRACT

The traditional exploitation of *Cistus ladanifer* L. (Cistaceae) species such as medicinal and aromatic plant (MAP) was studied for show the biodiversity changes associated to harvest and posterior regeneration. The results show significant increase of the species richness in all conditions with special high first year after harvest (>31 units), and in the Poaceae and Fabaceae families. Also, the geophytes habit increases and the reproductive capacities of the species. Globally the MAP exploitation of *C. ladanifer* is beneficial for the biodiversity of ecosystem.

Key words: Cistus, Richness, Biodiversity, Aromatic, Medicinal exploitation.

INTRODUCTION

Medicinal, aromatic and condimental species (MAP) are a constant source of consumption in indigenous populations and in rural areas far from large cities. Much of the origin of these natural products comes from wild collections in areas close to human settlements. In addition, they come from unique species that are often threatened and limited in their conservation, with plant species lost in some territories as a result of massive collections due to medicinal use and consumption (e.g. *Artemisia granatensis* Boiss (Blanca & Valle (1991), *Santolina elegans* Boiss. ex DC. (Blanca & al., 1990)).

Not all species exploited for medicinal purposes of wild origin are threatened, and their collection is a danger to the stability of the species or even to the habitat it occupies as can occur with species such as gentian, arnica, or mountain chamomile in the mountains of the Iberian Peninsula (Moreno & al., 2019).

Cistus ladanifer L. is a specie of huge applications in the pharmaceutical, cosmetic and perfumery industries. The origin of the production of *C. ladanifer* destined for industry in all cases is wild. In the last decade the productions of *C. ladanifer*, in the Mediterranean basin has been c. 10 t of essential oil (Pérez, 2022; Lubber & Verpoonte, 2011). The use of the rockrose in Iberian Peninsula dates to the prehistoric era (Buxó, 1993; 1997; Peña-Chocarro, 1995), the Neolithic settlers used the seeds of *C. ladanifer* in their domestic economy, being used to date (Frazao & al., 2022). Although the testimonies of its medicinal use in the Iberian Peninsula appear in the 16th century (Laguna, 1566), when it describes the process of extraction of the labdanum gum in the Sierra de Guadarrama mountains (Spain).

The manual or mechanized harvesting of *C. ladanifer* L. implies an increase in resources of great interest to the rural areas where is developed, together with an elimination of active biomass that can encourage natural fires in the Mediterranean forest and scrub. In addition, these collections are systematically been carried out on an annual basis in many locations of the SW of the Iberian

Península, promoting the over-exploitation of the natural cover of the scrubland and the potential losses of biodiversity in the territory, as has been shown in other areas where natural resources were overexploited.

The aim of this contribution is to facilitate an approximation of knowledge on changes in biodiversity in the wild populations of *C. ladanifer* L., which are traditionally harvested for exploitation as a medicinal, cosmetic or perfume plant.

DATA OF APPROACH TO STUDY

The area occupied by *C. ladanifer* populations susceptible to MAP harvesting in the countries where it is distributed (Algeria, France, Morocco, Portugal, Spain) (POWO, 2022) can reach more than 5 million ha, although the area traditionally used annually in wild collections does not exceed 3500 ha, mainly (>70 %) in the southern half of the border strip between Portugal and Spain. Harvesting takes place between July and September, being collected annually in Spain between 10000-20000 tonnes (Mediavilla & al., 2021) with a performance in essential oil ranging from 0.1-0.15%.

In addition, there is a use of the labdanum gum, obtained by immersion of the leaves and stems in warm carbonated water (55-60°C) followed by acidification of the medium (Frija & Afonso, 2014).

The harvesting is carried out in the upper third/half of the plants, always leaving a rest of the plants on the ground. Harvesting actions facilitate a loss of cover and shading capacity of the rockrose on the ground ranging from 30-65 %, increasing the degree of illumination of the soil surface. Along with the increase in the illuminated surface, the collections affect the survival of the plants, promoting plant losses between 10-22%. *C. ladanifer*, has allelopathic compounds in the leaves that inhibit the growth of vegetables, functioning as natural herbicides

(Chaves & Escudero, 1997; Alia & al., 2006, 2020; Chaves & al., 2019; Tena & al., 2021).

With the information provided we can understand the phenomena that occur on changes in biodiversity on the populations of *C. ladanifer*, used to obtain products intended for the pharmaceutical, cosmetic and perfume industries (essential oil and labdanum).

CHANGES IN THE BIODIVERSITY OF THE EXPLOITED POPULATIONS OF *CISTUS LADANIFER* L.

The populations studied during the years 2019-2020-2021 are located in the municipalities of Aliseda (Spain) (A), Barrado (Portugal) (B), Penamacor (Portugal) (P) and Villanueva del Fresno (Spain) (Vf). In all cases the upper fraction of the masses of *C. ladanifer* have been harvested during the months of July to August, leaving the lower fraction of the plants rooted in the ground. Data collection was carried out during the months of May and June looking for plant species that grow in the masses of *C. ladanifer*, prior to harvest, and then the following year after harvesting in the same period. In addition, data have been collected on plant species appearing in adjacent areas with poor *C. ladanifer* presence (<10% cover).

In all cases and situations studied, we have proceeded to evaluate the species richness based on the estimator for knowledge of richness in populations on the basis of capture and recapture of data, support on the works: Harris, 1959, and Chao, 1984, improved by Chao (1987) (Colwell & Coddington, 1996), supported by the collection of information from Ishii & al (2004), with an average number of samples in each case and period of X= 44-53 samples. Statistical analysis of means between the different population structures of *C. ladanifer* (Not harvest (J100); Harvest after 1 year (J50-1); Harvest after 2 years (J50-2); Poor *C. ladanifer* presence (NJ)) with a not parametric analysis Kruskal-Wallis test (Sokal & Rohlf, 1995).

Globally, 81 different plant species have been detected in the monitoring zones and periods, dominating the annual species of therophytes, typical of open spaces and poor soils in the SW half of the Iberian Peninsula (Table 1) (Rivas-Martinez, 1979).

The results provide an overview of the effect of the wild collections of *C. ladanifer* in the SW of Iberian Peninsula, where it has a flora-specific richness ($S2^*$) ranging from $S2^* = 5.1-12.3$ for homogeneous unexploited (collected) populations, up to $S2^* > 31$ in the same populations after one year of exploitation. After two years of exploitation the estimated richness decreases significantly, although it does not vary significantly from the increase observed in the first year (Figure 2). However, in cases where *C. ladanifer* has been reduced wealth increases considerably reached $S2^* = 49.3-65.1$ units (Figure 1).

The exploitation of *C. ladanifer* as MAP represents a reduction of at least 20 % in the leaf-producing fraction (Pérez, 2022) and allelopathic substances inhibiting the growth of accompanying vegetation (Alia & al., 2020; Chaves & al., 2019), especially herbaceous substances. After the collection of the upper fraction of the *C. ladanifer* specimens there is an increase in the insolation of the soil (Fernández-Mazuecos & Vargas, 2010; Frazao & al., 2018; Moreno & al., 2011; Valvueda & al., 1992), which may represent an increase of more than 50% compared to the time before the harvest. In addition, there is an increase in the rate of water from the rain that reaches the areas of the soil surface where a fraction of the allelopathic principles produced by *C. ladanifer* (Alia & al., 2006; Tena & al., 2021) are washed. The sum of these last two effects activates the sleeping seed bank and facilitates a more vigorous growth of the flora, increasing the specific richness and improving the biodiversity of the environment.

It should be noted that the presence of species of the Poaceae families (23.46 % representation), Fabaceae (16.50 % representation) and Asteraceae (11.00 % representation) (Table 1). Additionally, geophytes are observed in flowering of the families Orchidaceae (3.70 % representation) and Liliaceae (4.94 % representation), which usually only presented leaves when the masses of *C.*

ladanifer covered more than 80 % of the soil cover. The increase in specific richness, together with the emergence of reproductive structures in geophytes contribute even more to the increase in the global biodiversity of the environment.

Based on the follow up findings of the exploitation of *C. ladanifer*, as a MAP species in the SW of the Iberian Peninsula, it can be stated that there are positive elements in total biodiversity increased of the exploited areas and the removal of biomass (Ablanque & al., 2020) in these ecosystems with high risk of fires (Moreno & Oechel, 1994). The regular harvesting directly promotes the conservation of biodiversity: increasing specific richness, facilitating reproduction in plant species; increasing the rate of generative propagules in the seed bank and contributing to the promotion of other living beings dependent on plant diversity.




Figure 1. Relative abundance richness in four *Cistus ladanifer* population exploitation origins. Populations Alis: Aliseda (Spain); Barr: Barrado (Portugal); Pen: Penamacor (Portugal); Vfres: Villanueva del Fresno (Spain). Exploitation J100: Not harvest; J50-1: Harvest after 1 year; J50-2: Harvest after 2 years; NJ: Poor *Cistus ladanifer* presence.




Figure 2. Relative abundance richness in four *Cistus ladanifer* population structure types. J100: Not harvest; J50-1: Harvest after 1 year; J50-2: Harvest after 2 years; NJ: Poor *Cistus ladanifer* presence. **a, b, c;** indicate significant differences between samples ($p < 0,001$) (Kruskal Wallis test)

BIBLIOGRAPHY

Ablanque M, Vicente P, Cano C, Bados R & Plaza J (2020) Use of the essential oil of jara pringosa in the Sierra Norte from Madrid. 24th International Congress on Project Management and Engineering, 12 pp.

Alia JC, Sosa T, Escudero JC & Chaves N (2006) Autotoxicity Against Germination and Seedling Emergence in *Cistus ladanifer* L. *Plant and Soil* 282, 1, 327-332. DOI: 10.1007/s11104-005-6066-y

Alia JC, Caro JG, Campos VH & Chaves N (2020) Effect of Leaf Litter from *Cistus ladanifer* L. on the Germination and Growth of Accompanying Shrubland Species. *Plants*, 9, 593. <https://doi.org/10.3390/plants9050593>

Blanca G & Valle F (1991) Las plantas endémicas: *Artemisia granatensis*. Monograf. Fl. Veg. Béticas 6.

Blanca G, Valle F & Diaz de la Guardia C (1990) Las plantas endémicas de Andalucía oriental III. Monograf. Fl. Veg. Béticas 4/5, 3-44.

Bolaños MM & Lopez EG (1949) Jarales y jaras (Cistografía hispánica). Ediciones Ares. Madrid.

Buxó R (1993) Des semences et des fruits. Cueillette et Agriculture en France et en Espagne Mediterraneennes du Neolithique à l'Age du Fer. Phd. Montpellier University.

Buxó R (1997) Arqueología de las plantas. Ed. Crítica. Barcelona 367 pp

Chao A (1984) Non-parametric estimation of the number of classes in a population. *Scand. J. Stat.* 11, 265-270.

Chao A (1987) Estimating the population size for capture-recapture data with unequal catchability. *Biometrics* 43, 783-791.

Chaves N & Escudero JC (1997) Allelopathic effect of *Cistus ladanifer* on seed germination. *Functional Ecology*, 11, 4, 432-440. <https://doi.org/10.1046/j.1365-2435.1997.00107.x>

Chaves N, Ferrer I & Alías JC (2019). Autotoxicity of Diterpenes Present in Leaves of *Cistus ladanifer* L. *Plants* (Basel, Switzerland), 8(2), 27. <https://doi.org/10.3390/plants8020027>

Colwell RK & Coddington JA (1994) Estimating Terrestrial Biodiversity through Extrapolation. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 345, 101-118. <https://doi.org/10.1098/rstb.1994.0091>

Fernández-Mazuecos M & Vargas P (2010) Ecological rather than geographical isolation dominates quaternary formation of Mediterranean *Cistus* species. *Mol Ecol* 19(7):1381–1395. <https://doi.org/10.1111/j.1365-294X.2010.04549.x>

Frazão DF, Paulo L, Peres F, Resende M, Espírito Santo C, Barroca C, Moitinho A & Delgado F. (2022). *Cistus ladanifer* seeds: From ancient snack to novel and sustainable food ingredient. *Journal of Food Composition and Analysis*. 109. 104503. <https://doi.org/10.1016/j.jfca.2022.104503>

Frazão DF, Raimundo JR, Domingues JL, Quintela-Sabarís C, Gonçalves JC, Delgado F. (2018). *Cistus ladanifer* (Cistaceae): a natural resource in Mediterranean-type ecosystems. *Planta*, 247(2):289-300. <https://doi.org/10.1007/s00425-017-2825-2>

Frija LMT & Afonso CAM (2014) Supported acid-catalyzed flash vacuum thermolysis on the valorisation of labdanum resin. *Biomass and Bioenergy*, 71: 363-369. <https://doi.org/10.1016/j.biombioe.2014.09.020>

Harris B (1959) Determining the bounds on integrals with applications to cataloging problems. *Ann.Math. Statist.* 30, 521-548.

Ishii HT, Tanabe SI & Hiura T (2004) Exploring the Relationships Among Canopy Structure, Stand Productivity, and Biodiversity of Temperate Forest Ecosystems. *Forest Science*, 50, 3, 342–355. <https://doi.org/10.1093/forestscience/50.3.342>

Laguna A (1566). *Discorides*. Pedacio Dioscorides Anazarbeo, acerca de la Materia Medica Medicinal y de los Veneno Mortiferos. Salamanca. 616 pp.

Lubbe A & Verpoorte R (2011) Cultivation of Medicinal and Aromatic Plants for Specialty Industrial Materials. *Industrial Crops and Products*, 34, 785-801. <http://dx.doi.org/10.1016/j.indcrop.2011.01.019>

Mediavilla I, Blázquez MA, Ruiz A & Esteban LS (2021) Influence of the Storage of *Cistus ladanifer* L. Bales from Mechanised Harvesting on the Essential Oil Yield and Qualitative Composition. *Molecules* 26, 2379. <https://doi.org/10.3390/molecules26082379>

Moreno Saiz JC, Iriondo Alegría JM, Martínez García F, Martínez Rodríguez J & Salazar Mendías C (2019) *Atlas y Libro Rojo de la Flora Vasculár Amenazada de España*. Adenda 2017. Ministerio para la Transición Ecológica-Sociedad Española de Biología de la Conservación de Plantas. Madrid. 220 pp.

Moreno JM & Oechel WC (1994). *The role of fire in Mediterranean-Type Ecosystems*. *Ecological Studies* (Springer-Verlag) 107, 201 pp

Moreno JM, Zuazua E, Pérez B, Luna B, Velasco A & Dios VR (2011). Rainfall patterns after fire differentially affect the recruitment of three Mediterranean shrubs. *Biogeosciences* 8:3721–3732. <https://doi.org/10.5194/bg-8-3721-2011>

Peña-Chocarro L (1995) Prehistoric agricultura in southern Spain during the Neolithic and the Bronze Age: the application of ethnographic models. PhD College University London

Pérez C (2022) Bases ecologicas para la producción de aceite esencial de jara pringosa (*Cistus ladanifer* L.). PhD dissertation UEX. Plasencia (Spain). 235 pp.

POWO (2022) *Cistus ladanifer* L., taxonomy, distribution and bibliography. <https://powo.science.kew.org/taxon/168325-1>

Rivas-Martínez S (1979) Brezales y jarales de Europa occidental (Revisión fitosociológica de las clases Calluno-Ulicetea y Cisto-Lavanduletea). *Lazaroa*, 1, 5–127

Sokal RR & Rohlf FJ (1995) *Biometry. The principles and practice of statistics in biological research.* WH Freeman & Company. New York. 887 pp.

Tena C, Santiago ADR, Osuna D, Sosa T (2021) Phytotoxic Activity of p-Cresol, 2-Phenylethanol and 3-Phenyl-1-Propanol, Phenolic Compounds Present in *Cistus ladanifer* L. *Plants* (Basel) 10(6):1136. <https://doi.org/10.3390/plants10061136>.

Valbuena L, Tarrega R & Luis E (1992) Influence of heat on seed germination of *Cistus laurifolius* and *Cistus ladanifer*. *Int J Wildland Fire* 2(1):15–20. <https://doi.org/10.1071/WF9920015>